

Supplementary Derivations for Nonparametric Stochastic Discount Factor Decomposition

Timothy M. Christensen

April 5, 2017

This note works through parametric derivations required for the simulations and empirical application in the paper “Nonparametric Stochastic Discount Factor Decomposition”.

1 Recursive preferences with linear-Gaussian state

1.1 Univariate case

First assume that:

$$g_{t+1} - \mu = \kappa(g_t - \mu) + \sigma e_{t+1}, \quad e_{t+1} \sim \text{i.i.d. } N(0, 1). \quad (1)$$

Step 1: fixed-point problem.

Guess a solution of the form $h(g) = e^{b_0 + b_1 g}$ to the fixed-point equation:

$$h(g_t) = \mathbb{E}[G_{t+1}^{1-\gamma} (h(g_{t+1}))^\beta | g_t]. \quad (2)$$

Then we have:

$$e^{b_0 + b_1 g_t} = \mathbb{E}[e^{(1-\gamma)g_{t+1} + \beta b_0 + \beta b_1 g_{t+1}} | g_t] \quad (3)$$

$$= e^{\beta b_0} \mathbb{E}[e^{(1-\gamma + \beta b_1)g_{t+1}} | g_t] \quad (4)$$

$$= e^{\beta b_0} e^{(1-\gamma + \beta b_1)(1-\kappa)\mu + (1-\gamma + \beta b_1)\kappa g_t + \frac{1}{2}(1-\gamma + \beta b_1)^2 \sigma^2}. \quad (5)$$

Collecting coefficients of g_t :

$$b_1 = \kappa(1 - \gamma + \beta b_1) \quad \Rightarrow \quad b_1 = \frac{\kappa(1 - \gamma)}{1 - \kappa\beta}. \quad (6)$$

Collecting constants:

$$b_0 = \beta b_0 + (1 - \gamma + \beta b_1)(1 - \kappa)\mu + \frac{1}{2}(1 - \gamma + \beta b_1)^2 \sigma^2 \quad (7)$$

$$\Rightarrow b_0 = \frac{1}{1 - \beta} \left(b_2(1 - \kappa)\mu + \frac{1}{2} b_2^2 \sigma^2 \right) \quad (8)$$

with $b_2 = 1 - \gamma + \beta b_1$.

Also notice that unconditionally $g_t \sim N(\mu, \frac{\sigma^2}{1 - \kappa^2})$. Therefore:

$$\|h\|^2 = e^{2b_0} \mathbb{E}[e^{2b_1 g_t}] = e^{2b_0 + 2b_1 \mu + 2b_1^2 \frac{\sigma^2}{1 - \kappa^2}} \Rightarrow \|h\| = e^{b_0 + b_1 \mu + b_1^2 \frac{\sigma^2}{1 - \kappa^2}} \quad (9)$$

and so:

$$\chi(g_t) = \frac{e^{b_0 + b_1 g_t}}{e^{b_0 + b_1 \mu + b_1^2 \frac{\sigma^2}{1 - \kappa^2}}} = e^{-(b_1 \mu + b_1^2 \frac{\sigma^2}{1 - \kappa^2}) + b_1 g_t} = e^{c_0 + b_1 g_t} \quad (10)$$

where $c_0 = -(b_1 \mu + b_1^2 \frac{\sigma^2}{1 - \kappa^2})$. Finally,

$$\lambda = e^{(1 - \beta)(b_0 + b_1 \mu + b_1^2 \frac{\sigma^2}{1 - \kappa^2})} = e^{(1 - \beta)(b_0 - c_0)} . \quad (11)$$

Step 2: Perron-Frobenius problem.

First, write the SDF as:

$$m(g_t, g_{t+1}) = \frac{\beta}{\lambda} G_{t+1}^{-\gamma} \frac{\chi(g_{t+1})^\beta}{\chi(g_t)} \quad (12)$$

$$= \frac{\beta}{\lambda} e^{-\gamma g_{t+1} + \beta(c_0 + b_1 g_{t+1}) - (c_0 + b_1 g_t)} \quad (13)$$

$$= \frac{\beta}{\lambda} e^{(\beta - 1)c_0} e^{(\beta b_1 - \gamma)g_{t+1} - b_1 g_t} . \quad (14)$$

This gives:

$$\mathbb{E}[\log m(g_t, g_{t+1})] = \log \left(\frac{\beta}{\lambda} \right) + (\beta - 1)c_0 + (\beta b_1 - \gamma - b_1)\mu \quad (15)$$

which is needed for the entropy calculation.

Guess a solution of the form $\phi(g) = e^{a_0 + a_1 g}$ and plugging into $\mathbb{M}\phi = \rho\phi$, we obtain:

$$\rho e^{a_1 g_t} = \frac{\beta}{\lambda} e^{(\beta - 1)c_0} e^{-b_1 g_t} \mathbb{E}[e^{(a_1 + \beta b_1 - \gamma)g_{t+1}} | g_t] \quad (16)$$

$$= \frac{\beta}{\lambda} e^{(\beta - 1)c_0} e^{-b_1 g_t} e^{(a_1 + \beta b_1 - \gamma)(1 - \kappa)\mu + (a_1 + \beta b_1 - \gamma)\kappa g_t + \frac{1}{2}(a_1 + \beta b_1 - \gamma)^2 \sigma^2} . \quad (17)$$

Collecting coefficients of g_t :

$$a_1 = -b_1 + (a_1 + \beta b_1 - \gamma)\kappa \quad \Rightarrow \quad a_1 = \frac{-b_1 + \kappa(\beta b_1 - \gamma)}{1 - \kappa} \quad (18)$$

and so $a_0 = -a_1\mu - a_1^2 \frac{\sigma^2}{1-\kappa^2}$.

Moreover,

$$\rho = \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{(a_1+\beta b_1-\gamma)(1-\kappa)\mu + \frac{1}{2}(a_1+\beta b_1-\gamma)^2\sigma^2} = \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{\nu(1-\kappa)\mu + \frac{1}{2}\nu^2\sigma^2} \quad (19)$$

where $\nu = \beta b_1 - \gamma + a_1$.

Step 2b: time-reversed Perron-Frobenius problem.

Now we guess a solution of the form $\phi(g) = e^{a_0^* + a_1^* g}$. Plugging into $\mathbb{M}^* \phi^* = \rho \phi^*$ and using reversibility of $\{g_t\}$ (Weiss, 1975), we obtain:

$$\rho e^{a_1^* g_{t+1}} = \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{(\beta b_1 - \gamma)g_{t+1}} \mathbb{E}[e^{(a_1^* - b_1)g_t} | g_{t+1}] \quad (20)$$

$$= \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{(\beta b_1 - \gamma)g_{t+1} + (a_1^* - b_1)(1-\kappa)\mu + (a_1^* - b_1)\kappa g_{t+1} + \frac{1}{2}(a_1^* - b_1)^2\sigma^2}. \quad (21)$$

Collecting coefficients of g_{t+1} :

$$a_1^* = \beta b_1 - \gamma + (a_1^* - b_1)\kappa \quad \Rightarrow \quad a_1^* = \frac{(\beta - \kappa)b_1 - \gamma}{1 - \kappa}. \quad (22)$$

We choose a_0^* so that:

$$1 = \mathbb{E}[e^{a_0 + a_0^* + (a_1 + a_1^*)g_t}] = e^{a_0 + a_0^* + (a_1 + a_1^*)\mu + \frac{1}{2}(a_1 + a_1^*)^2 \frac{\sigma^2}{1-\kappa^2}} \quad (23)$$

which implies $a_0^* = -(a_0 + (a_1 + a_1^*)\mu + \frac{1}{2}(a_1 + a_1^*)^2 \frac{\sigma^2}{1-\kappa^2})$.

Step 3: Covariance of the permanent and transitory components.

The permanent component is:

$$\frac{M_{t+1}^P}{M_t^P} = \frac{\frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{(\beta b_1 - \gamma)g_{t+1} - b_1 g_t}}{\frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{\nu(1-\kappa)\mu + \frac{1}{2}\nu^2\sigma^2}} e^{a_1 g_{t+1} - a_1 g_t} = e^{-\nu(1-\kappa)\mu - \frac{1}{2}\nu^2\sigma^2} e^{\nu g_{t+1} - (b_1 + a_1)g_t}. \quad (24)$$

Therefore:

$$m_{t+1}^P = \log\left(\frac{M_{t+1}^P}{M_t^P}\right) = \text{const} - (b_1 + a_1)g_t + \nu g_{t+1}. \quad (25)$$

The transitory component is:

$$\frac{M_{t+1}^T}{M_t^T} = \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{\nu(1-\kappa)\mu + \frac{1}{2}\nu^2\sigma^2} e^{a_1g_t - a_1g_{t+1}} \quad (26)$$

and so:

$$m_{t+1}^T = \log\left(\frac{M_{t+1}^T}{M_t^T}\right) = \text{const} + a_1g_t - a_1g_{t+1}. \quad (27)$$

Notice that:

$$\begin{pmatrix} g_t \\ g_{t+1} \end{pmatrix} \sim N\left(\begin{pmatrix} \mu \\ \mu \end{pmatrix}, \underbrace{\begin{pmatrix} \frac{\sigma^2}{1-\kappa^2} & \kappa\frac{\sigma^2}{1-\kappa^2} \\ \kappa\frac{\sigma^2}{1-\kappa^2} & \frac{\sigma^2}{1-\kappa^2} \end{pmatrix}}_{\mathbb{V}}\right). \quad (28)$$

Therefore:

$$\text{Cov}(m_{t+1}^P, m_{t+1}^T) = v_P' \mathbb{V} v_T \quad \text{Var}(m_{t+1}^P) = v_P' \mathbb{V} v_P \quad \text{Var}(m_{t+1}^T) = v_T' \mathbb{V} v_T \quad (29)$$

where:

$$v_P = \begin{pmatrix} -(a_1 + b_1) \\ \nu \end{pmatrix} \quad v_T = \begin{pmatrix} a_1 \\ -a_1 \end{pmatrix}. \quad (30)$$

1.2 Multivariate case

Now assume that $g_{t+1} = \Gamma' X_{t+1}$ where

$$X_{t+1} - \mu = A(X_t - \mu) + \sigma e_{t+1}, \quad e_{t+1} \sim \text{i.i.d. N}(0, I) \quad (31)$$

and let $\Sigma = \sigma\sigma'$.

Step 1: fixed-point problem.

Guess a solution of the form $h(x) = e^{b_0 + b_1'x}$ to the fixed-point equation:

$$h(X_t) = \mathbb{E}[G_{t+1}^{1-\gamma} (h(X_{t+1}))^\beta | X_t] \quad (32)$$

$$\Rightarrow e^{b_0 + b_1'X_t} = \mathbb{E}[e^{(1-\gamma)\Gamma'X_{t+1} + \beta b_0 + \beta b_1'X_{t+1}} | X_t] \quad (33)$$

$$= e^{\beta b_0} \mathbb{E}[e^{((1-\gamma)\Gamma + \beta b_1)'X_{t+1}} | X_t] \quad (34)$$

$$= e^{\beta b_0} e^{((1-\gamma)\Gamma + \beta b_1)'(I-A)\mu + ((1-\gamma)\Gamma + \beta b_1)'AX_t + \frac{1}{2}((1-\gamma)\Gamma + \beta b_1)'\Sigma((1-\gamma)\Gamma + \beta b_1)}. \quad (35)$$

Collecting coefficients of X_t :

$$b_1 = A'((1 - \gamma)\Gamma + \beta b_1) \quad \Rightarrow \quad b_1 = (1 - \gamma)(I - \beta A')^{-1} A' \Gamma. \quad (36)$$

Collecting constants:

$$b_0 = \beta b_0 + ((1 - \gamma)\Gamma + \beta b_1)'(I - A)\mu + \frac{1}{2}((1 - \gamma)\Gamma + \beta b_1)'\Sigma((1 - \gamma)\Gamma + \beta b_1) \quad (37)$$

$$\Rightarrow b_0 = \frac{1}{1 - \beta} \left(b_2'(I - A)\mu + \frac{1}{2} b_2'\Sigma b_2 \right) \quad (38)$$

where $b_2 = \beta b_1 + (1 - \gamma)\Gamma$.

The marginal of X_t is $N(\mu, V)$ where V solves $V = AVA' + \Sigma$. Therefore:

$$\|h\|^2 = e^{2b_0} \mathbb{E}[e^{2b_1' X_t}] = e^{2b_0 + 2b_1' \mu + 2b_1' V b_1} \quad \Rightarrow \quad \|h\| = e^{b_0 + b_1' \mu + b_1' V b_1} \quad (39)$$

and so

$$\chi(X_t) = \frac{e^{b_0 + b_1' x}}{e^{b_0 + b_1' \mu + b_1' V b_1}} = e^{-(b_1' \mu + b_1' V b_1) + b_1' X_t} = e^{c_0 + b_1' X_t} \quad (40)$$

where $c_0 = -(b_1' \mu + b_1' V b_1)$. Finally,

$$\lambda = e^{(1 - \beta)(b_0 + b_1' \mu + b_1' V b_1)} = e^{(1 - \beta)(b_0 - c_0)}. \quad (41)$$

Step 2: Perron-Frobenius problem.

First, write the SDF as:

$$m(X_t, X_{t+1}) = \frac{\beta}{\lambda} G_{t+1}^{-\gamma} \frac{\chi(X_{t+1})^\beta}{\chi(X_t)} \quad (42)$$

$$= \frac{\beta}{\lambda} e^{-\gamma \Gamma' X_{t+1} + \beta(c_0 + b_1' X_{t+1}) - (c_0 + b_1' X_t)} \quad (43)$$

$$= \frac{\beta}{\lambda} e^{(\beta - 1)c_0} e^{(\beta b_1 - \gamma \Gamma)' X_{t+1} - b_1' X_t}. \quad (44)$$

This gives:

$$\mathbb{E}[\log m(X_t, X_{t+1})] = \log \left(\frac{\beta}{\lambda} \right) + (\beta - 1)c_0 + (\beta b_1 - \gamma \Gamma - b_1)' \mu \quad (45)$$

which is needed for the entropy calculation.

Guess a solution of the form $\phi(x) = e^{a_0 + a_1'x}$. Plugging into $M\phi = \rho\phi$, we obtain:

$$\rho e^{a_1'X_t} = \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{-b_1'X_t} \mathbb{E}[e^{(a_1 + \beta b_1 - \gamma\Gamma)'X_{t+1}} | X_t] \quad (46)$$

$$= \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{-b_1'X_t} e^{(a_1 + \beta b_1 - \gamma\Gamma)'(I-A)\mu + (a_1 + \beta b_1 - \gamma\Gamma)'AX_t + \frac{1}{2}(a_1 + \beta b_1 - \gamma\Gamma)'\Sigma(a_1 + \beta b_1 - \gamma\Gamma)}. \quad (47)$$

Collecting coefficients of X_t :

$$a_1 = -b_1 + A'(a_1 + \beta b_1 - \gamma\Gamma) \quad \Rightarrow \quad a_1 = (I - A')^{-1}(-b_1 + A'(\beta b_1 - \gamma\Gamma)). \quad (48)$$

Moreover,

$$\rho = \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{(a_1 + \beta b_1 - \gamma\Gamma)'(I-A)\mu + \frac{1}{2}(a_1 + \beta b_1 - \gamma\Gamma)'\Sigma(a_1 + \beta b_1 - \gamma\Gamma)} = \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{\nu'(I-A)\mu + \frac{1}{2}\nu'\Sigma\nu} \quad (49)$$

where $\nu = \beta b_1 - \gamma\Gamma + a_1$.

Step 3: Covariance of the permanent and transitory components.

The permanent component is:

$$\frac{M_{t+1}^P}{M_t^P} = \frac{\frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{(\beta b_1 - \gamma\Gamma)'X_{t+1} - b_1'X_t}}{\frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{\nu'(I-A)\mu + \frac{1}{2}\nu'\Sigma\nu}} e^{a_1'X_{t+1} - a_1'X_t} = e^{-\nu'(I-A)\mu - \frac{1}{2}\nu'\Sigma\nu} e^{\nu'X_{t+1} - (a_1 + b_1)'X_t} \quad (50)$$

and so:

$$m_{t+1}^P = \text{const} - (a_1 + b_1)'X_t + \nu'X_{t+1}. \quad (51)$$

The transitory component is:

$$\frac{M_{t+1}^T}{M_t^T} = \frac{\beta}{\lambda} e^{(\beta-1)c_0} e^{\nu'(I-A)\mu + \frac{1}{2}\nu'\Sigma\nu} e^{a_1'X_t - a_1'X_{t+1}} \quad (52)$$

and so:

$$m_{t+1}^T = \text{const} + a_1'X_t - a_1'X_{t+1}. \quad (53)$$

Notice that:

$$\begin{pmatrix} X_t \\ X_{t+1} \end{pmatrix} \sim N\left(\begin{pmatrix} \mu \\ \mu \end{pmatrix}, \underbrace{\begin{pmatrix} V & VA' \\ AV & V \end{pmatrix}}_V\right). \quad (54)$$

Therefore, the covariance of m_{t+1}^P and m_{t+1}^T are as in display (29) with \mathbb{V} from (54) and:

$$v_P = \begin{pmatrix} -(a_1 + b_1) \\ \nu \end{pmatrix} \quad v_T = \begin{pmatrix} a_1 \\ -a_1 \end{pmatrix}. \quad (55)$$

2 Recursive preferences with stochastic volatility in consumption growth

2.1 Derivation

We use a specification of the consumption growth process similar to one studied in [Backus, Chernov, and Zin \(2014\)](#) but without habits. Assume that:

$$\begin{aligned} g_{t+1} - \mu &= \kappa(g_t - \mu) + v_t^{1/2} e_{t+1}, \quad e_{t+1} \sim \text{i.i.d. } N(0, 1) \\ v_{t+1} &= \text{ARG}(c_v, \varphi_v, \delta_v) \end{aligned}$$

where $\text{ARG}(c_v, \varphi_v, \delta_v)$ is an autoregressive gamma (ARG) process of order 1 parameterized by $(c_v, \varphi_v, \delta_v)$ that is independent of the sequence of $N(0, 1)$ innovations. We refer the reader to [Gourieroux and Jasiak \(2006\)](#) for background material on ARG processes.

The state vector is $X_t = (g_t, v_t)'$. The following derivations are essentially a special case of Appendix H in [Backus et al. \(2014\)](#) but are included for completeness. Throughout, we will use the fact that:

$$\mathbb{E}[e^{sv_{t+1}} | v_t] = e^{\frac{\varphi_v s}{1-sc_v} v_t - \delta_v \log(1-sc_v)} \quad (56)$$

(see Appendix H in [Backus et al. \(2014\)](#)).

Step 1: fixed-point problem.

Guess a solution of the form $h(g, v) = e^{a_0 + b_1 g + a_2 v}$ to the fixed-point equation:

$$h(g_t, v_t) = \mathbb{E}[G_{t+1}^{1-\gamma} (h(g_{t+1}, v_{t+1}))^\beta | g_t, v_t] \quad (57)$$

$$\Rightarrow e^{a_0 + a_1 g_t + a_2 v_t} = \mathbb{E}[e^{(1-\gamma)g_{t+1} + \beta a_0 + \beta a_1 g_{t+1} + \beta a_2 v_{t+1}} | g_t, v_t] \quad (58)$$

$$= e^{\beta a_0} \mathbb{E}[e^{(1-\gamma + \beta a_1)g_{t+1} + \beta a_2 v_{t+1}} | g_t, v_t] \quad (59)$$

$$= e^{\beta a_0} e^{(1-\gamma + \beta a_1)(1-\kappa)\mu + (1-\gamma + \beta a_1)\kappa g_t + \frac{1}{2}(1-\gamma + \beta a_1)^2 v_t} \mathbb{E}[e^{\beta a_2 v_{t+1}} | v_t] \quad (60)$$

$$= e^{\beta a_0} e^{(1-\gamma + \beta a_1)(1-\kappa)\mu + (1-\gamma + \beta a_1)\kappa g_t + \frac{1}{2}(1-\gamma + \beta a_1)^2 v_t} e^{\frac{\varphi_v \beta a_2}{1-\beta a_2 c_v} v_t - \delta_v \log(1-\beta a_2 c_v)}. \quad (61)$$

Collecting coefficients of g_t :

$$a_1 = \kappa(1 - \gamma + \beta a_1) \quad \Rightarrow \quad a_1 = \frac{\kappa(1 - \gamma)}{1 - \beta\kappa}. \quad (62)$$

Collecting coefficients of v_t :

$$a_2 = \frac{1}{2}(1 - \gamma + \beta a_1)^2 + \frac{\varphi_v \beta a_2}{1 - \beta a_2 c_v} \quad \Rightarrow \quad a_2 - \nu = \frac{\varphi_v \beta a_2}{1 - \beta a_2 c_v} \quad (63)$$

where $\nu = \frac{1}{2}(1 - \gamma + \beta a_1)^2$. Rearranging:

$$0 = (a_2 - \nu)(1 - \beta c_v a_2) - \beta \varphi_v a_2 \quad (64)$$

$$= \beta c_v a_2^2 + (\beta(\varphi_v - c_v \nu) - 1)a_2 + \nu \quad (65)$$

so we solve the quadratic and take the smallest root to obtain:

$$a_2 = \frac{-(\beta(\varphi_v - c_v \nu) - 1) - \sqrt{(\beta(\varphi_v - c_v \nu) - 1)^2 - 4\beta c_v \nu}}{2\beta c_v} \quad (66)$$

provided $(\beta(\varphi_v - c_v \nu) - 1)^2 - 4\beta c_v \nu \geq 0$. Taking the smallest root ensures that $a_2 = 0$ if the variance of $\log g_t$ is zero (see Appendix H in [Backus et al. \(2014\)](#)).

Collecting the constant terms:

$$a_0 = \beta a_0 + (1 - \gamma + \beta a_1)(1 - \kappa)\mu - \delta_v \log(1 - \beta a_2 c_v) \quad (67)$$

$$\Rightarrow a_0 = \frac{1}{1 - \beta} ((1 - \gamma + \beta a_1)(1 - \kappa)\mu - \delta_v \log(1 - \beta a_2 c_v)) \quad (68)$$

provided $\beta a_2 c_v < 1$.

Step 2: Perron-Frobenius problem.

First, write the SDF as:

$$m(X_t, X_{t+1}) = \beta G_{t+1}^{-\gamma} \frac{h(X_{t+1})^\beta}{h(X_t)} \quad (69)$$

$$= \beta e^{-\gamma g_{t+1} + \beta(a_0 + a_1 g_{t+1} + a_2 v_{t+1}) - (a_0 + a_1 g_t + a_2 v_t)} \quad (70)$$

$$= \beta e^{(\beta-1)a_0 + (\beta a_1 - \gamma)g_{t+1} + \beta a_2 v_{t+1} - a_1 g_t - a_2 v_t} \quad (71)$$

$$= d_0 e^{d_1 g_{t+1} + d_2 v_{t+1} + d_3 g_t + d_4 v_t} \quad (72)$$

where

$$d_0 = \beta e^{(\beta-1)a_0} \quad d_1 = \beta a_1 - \gamma \quad d_2 = \beta a_2 \quad d_3 = -a_1 \quad d_4 = -a_2. \quad (73)$$

In this notation we have:

$$\mathbb{E}[\log m(X_t, X_{t+1})] = \log(d_0) + (d_1 + d_3)\mu + (d_2 + d_4) \frac{\delta_v c_v}{1 - \varphi_v} \quad (74)$$

which is needed for the entropy of the permanent component.

Conjecture a solution of the form $\phi(g, v) = e^{c_0 + c_1 g + c_2 v}$. Plugging into $\mathbb{M}\phi = \rho\phi$ gives:

$$\rho e^{c_1 g_t + c_2 v_t} = d_0 e^{d_3 g_t + d_4 v_t} \mathbb{E}[e^{(d_1 + c_1)g_{t+1} + (d_2 + c_2)v_{t+1}} | g_t, v_t] \quad (75)$$

$$= d_0 e^{d_3 g_t + d_4 v_t} e^{(d_1 + c_1)(1-\kappa)\mu + (d_1 + c_1)\kappa g_t + \frac{1}{2}(d_1 + c_1)^2 v_t} \mathbb{E}[e^{(d_2 + c_2)v_{t+1}} | v_t] \quad (76)$$

$$= d_0 e^{(d_1 + c_1)(1-\kappa)\mu} e^{[d_3 + (d_1 + c_1)\kappa]g_t + [d_4 + \frac{1}{2}(d_1 + c_1)^2]v_t} \mathbb{E}[e^{(d_2 + c_2)v_{t+1}} | v_t] \quad (77)$$

$$= d_0 e^{(d_1 + c_1)(1-\kappa)\mu} e^{[d_3 + (d_1 + c_1)\kappa]g_t + [d_4 + \frac{1}{2}(d_1 + c_1)^2]v_t} e^{\frac{\varphi_v(d_2 + c_2)}{1 - c_v(d_2 + c_2)} v_t - \delta_v \log(1 - c_v(d_2 + c_2))} \quad (78)$$

$$= d_0 e^{(d_1 + c_1)(1-\kappa)\mu - \delta_v \log(1 - c_v(d_2 + c_2))} e^{[d_3 + (d_1 + c_1)\kappa]g_t + [d_4 + \frac{1}{2}(d_1 + c_1)^2 + \frac{\varphi_v(d_2 + c_2)}{1 - c_v(d_2 + c_2)}]v_t}. \quad (79)$$

Collecting coefficients of g_t :

$$c_1 = d_3 + \kappa(d_1 + c_1) \quad \Rightarrow \quad c_1 = \frac{d_3 + \kappa d_1}{1 - \kappa}. \quad (80)$$

Collecting coefficients of v_t :

$$c_2 = d_4 + \frac{1}{2}(d_1 + c_1)^2 + \frac{\varphi_v(d_2 + c_2)}{1 - c_v(d_2 + c_2)}. \quad (81)$$

Let $c_3 = c_2 + d_2$. Then:

$$c_3 = d_2 + d_4 + \frac{1}{2}(d_1 + c_1)^2 + \frac{\varphi_v c_3}{1 - c_v c_3} = \xi + \frac{\varphi_v c_3}{1 - c_v c_3}. \quad (82)$$

where $\xi = d_2 + d_4 + \frac{1}{2}(d_1 + c_1)^2$. Rearranging:

$$(c_3 - \xi)(1 - c_v c_3) = \varphi_v c_3 \quad (83)$$

$$\Rightarrow 0 = c_v c_3^2 + (\varphi_v - 1 - c_v \xi)c_3 + \xi. \quad (84)$$

Solving the quadratic and again taking the negative root, we obtain:

$$c_3 = \frac{-(\varphi_v - 1 - c_v \xi) - \sqrt{(\varphi_v - 1 - c_v \xi)^2 - 4c_v \xi}}{2c_v} \quad (85)$$

provided $(\varphi_v - 1 - c_v \xi)^2 - 4c_v \xi \geq 0$. Finally, set $c_2 = c_3 - d_2$.

We also obtain

$$\rho = d_0 e^{(d_1 + c_1)(1 - \kappa)\mu - \delta_v \log(1 - c_v(d_2 + c_2))} \quad (86)$$

provided $c_v(d_2 + c_2) < 1$.

Step 3: Covariance of the permanent and transitory components.

The permanent component is:

$$\frac{M_{t+1}^P}{M_t^P} = \frac{d_0}{d_0 e^{(d_1 + c_1)(1 - \kappa)\mu - \delta_v \log(1 - c_v(d_2 + c_2))}} e^{(d_1 + c_1)g_{t+1} + (d_2 + c_2)v_{t+1} + (d_3 - c_1)g_t + (d_4 - c_2)v_t} \quad (87)$$

and so:

$$m_{t+1}^P = \text{const} + \begin{pmatrix} d_3 - c_1 \\ d_4 - c_2 \end{pmatrix}' X_t + \begin{pmatrix} d_1 + c_1 \\ d_2 + c_2 \end{pmatrix}' X_{t+1}. \quad (88)$$

The transitory component is:

$$\frac{M_{t+1}^T}{M_t^T} = d_0 e^{(d_1 + c_1)(1 - \kappa)\mu - \delta \log(1 - c_v(d_2 + c_2))} e^{-c_1 g_{t+1} - c_2 v_{t+1} + c_1 g_t + c_2 v_t} \quad (89)$$

and so:

$$m_{t+1}^T = \text{const} + \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}' X_t - \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}' X_{t+1}. \quad (90)$$

It remains to construct the covariance matrix \mathbb{V} of (X_t, X_{t+1}) . From [Gourieroux and Jasiak \(2006\)](#) we have:

$$\mathbb{E}[v_t] = \frac{c_v \delta_v}{1 - \varphi_v} \quad (91)$$

$$\text{Var}(v_t) = \frac{c_v^2 \delta_v}{(1 - \varphi_v)^2} \quad (92)$$

$$\text{Cov}(v_t, v_{t+1}) = \varphi_v \text{Var}(v_t) = \frac{\varphi_v c_v^2 \delta_v}{(1 - \varphi_v)^2}. \quad (93)$$

It follows that

$$\text{Var}(g_t) = \frac{c_v \delta_v}{(1 - \varphi_v)(1 - \kappa^2)} \quad (94)$$

$$\text{Cov}(g_t, g_{t+1}) = \kappa \text{Var}(g_t) = \frac{\kappa c_v \delta_v}{(1 - \varphi_v)(1 - \kappa^2)}. \quad (95)$$

Finally, note that $\text{Cov}(g_t, v_{t+1})$, $\text{Cov}(v_t, g_{t+1})$, and $\text{Cov}(v_{t+1}, g_{t+1})$ are all zero. The covariance matrix of $(X'_t, X'_{t+1})'$ is therefore:

$$\mathbb{V} = \begin{pmatrix} \frac{c_v \delta_v}{(1 - \varphi_v)(1 - \kappa^2)} & 0 & \kappa \frac{c_v \delta_v}{(1 - \varphi_v)(1 - \kappa^2)} & 0 \\ 0 & \frac{c_v^2 \delta_v}{(1 - \varphi_v)^2} & 0 & \varphi_v \frac{c_v^2 \delta_v}{(1 - \varphi_v)^2} \\ \kappa \frac{c_v \delta_v}{(1 - \varphi_v)(1 - \kappa^2)} & 0 & \frac{c_v \delta_v}{(1 - \varphi_v)(1 - \kappa^2)} & 0 \\ 0 & \varphi_v \frac{c_v^2 \delta_v}{(1 - \varphi_v)^2} & 0 & \frac{c_v^2 \delta_v}{(1 - \varphi_v)^2} \end{pmatrix}. \quad (96)$$

The covariance of m_{t+1}^P and m_{t+1}^T are as in display (29) with this \mathbb{V} and with:

$$v_P = \begin{pmatrix} d_3 - c_1 \\ d_4 - c_2 \\ d_1 + c_1 \\ d_2 + c_2 \end{pmatrix} \quad v_T = \begin{pmatrix} c_1 \\ c_2 \\ -c_1 \\ -c_2 \end{pmatrix}. \quad (97)$$

2.2 Estimation

We estimate the parameters $(\mu, \kappa, \varphi_v, c_v, \delta_v)$ by indirect inference. The auxiliary model for $\{g_t\}$ is an AR(1) process with a GARCH(1,1) process fit to the AR(1) residuals. We use 500 simulated series. In fitting the auxiliary model, we first estimate the AR(1) parameters μ and κ by regression then and fit the GARCH(1,1) process to the residuals by maximum likelihood.

References

- Backus, D., M. Chernov, and S. Zin (2014). Sources of entropy in representative agent models. *Journal of Finance* 69(1), 51–99.
- Gourieroux, C. and J. Jasiak (2006). Autoregressive gamma processes. *Journal of Forecasting* 25(2), 129–152.
- Weiss, G. (1975). Time-reversibility of linear stochastic processes. *Journal of Applied Probability* 12(4), 831–836.